## JC20 Rec'd PCT/PTO 0 4 MAR 2002

	PTO-139 11-2000		IMERCE PATENT AND TRADEMARK OFFICE	ATTORNEYS DOCKET NUMBER RN99110			
	TRA	ANSMITTAL LETTER T	O THE UNITED STATES	U.S. APPLICATION NO. (JF KNOWN)			
D			O/EO/US) CONCERNING A	10/070632			
		FILING UNDER	-	NOT APPLICABLE 010072			
		TIONAL APPLICATION NO. <b>00/02452</b>	INTERNATIONAL FILING DATE September 06, 2000	PRIORITY DATE CLAIMED September 07, 2000			
	TITLE OF INVENTION						
PR	OC	ESS FOR PRODUC	TION OF EXOPOLYSACO	CHARIDES			
		***************************************					
		NT (S) FOR DO/EO/US:	CIMON				
	vier	NORE and Jean-Luc	SIMON				
Appl	icant l	nerewith submits the United State	es Designated/Elected Office (DO/EO/US)	the following items and other information:			
1.	$\boxtimes$	This is a FIRST submission of	items concerning a filing under 35 U.S.C.	371.			
2.		This is a SECOND or SUBSEC	QUENT submission of items concerning a	filing under 35 U.S.C. 371.			
3.	$\boxtimes$		gin national examination procedures (35 U.	S.C. 371 (f)). The submission must include items (5), (6), (9)			
١,		and 24 indicated below.	association of 10 months from the majority	data (Autiala 21)			
4. 5.	M	•	expiration of 19 months from the priority of the expiration as filed (35 U.S.C. 371(c) (2))	nate (Atticle 31).			
J.	$\boxtimes$		required only if not communicated by the I	nternational Bureau)			
		_	cated by the International bureau.	mermaronai zareaa).			
		_	he application was filed in the United State	es Receiving Office (RO/US).			
6.	$\boxtimes$	-	of the International Application as filed (3				
		a. 🛛 is attached hereto.					
			y submitted under 35 U.S.C. 154(d)(4).				
7.	$\boxtimes$		ne International Application under PCT Art	icle 19 (35 U S C 371(c)(3))			
	_	a. are attached hereto	(required only if not communicated by the	International Bureau).			
		b. have been commur	nicated by the International Bureau.				
		c. have not been mad	e; however, the time limit for making such	amendments has NOT expired.			
		d. 🛛 have not been mad	e and will not be made.				
8.		An English language translation	of the amendments to the claims under PC	CT Article 19 (35 U S C 371(c)(3)).			
9.	$\boxtimes$	An oath or declaration of the in	ventor(s) (35 U.S.C. 371(c)(4)). (UNSIGN	ED)			
10.		An English language translation $371(c)(5)$ .	n of the annexes to the International Prelimi	inary Examination Report under PCT Article 36 (35 U.S.C.			
11.	$\boxtimes$		liminary Examination Report (PCT/IPEA/4	09).			
12.	$\boxtimes$	A copy of the International Sea	rch Report (PCT/ISA/210)				
Item	s 13 to	o 18 below concern document(s)	or information included:				
13.	$\boxtimes$	An information Disclosure State	ement under 37 CFR 1 97 and 1 98.				
14.		An assignment document for re	cording. A separate cover sheet in complian	nce with 37 CFR 3 28 and 3 31 is included.			
15.	$\boxtimes$	A FIRST preliminary amendme	ent.				
16		A SECOND or SUBSEQUENT	r preliminary amendment.				
17.		A substitute specification.					
18.		A change of power of attorney		D. 10. 0. 105 V.G.C. 1001 1005			
19.		·	ie sequence listing in accordance with PCT				
20.			international application under 35 U S.C.				
21. 22.			anguage translation of the international app	meation under 33 O.S.C. 134(d)(4).			
23.		Certificate of Mailing by Expre Other items or information:	55 I <b>VIAII</b> .				
22.			PCT/IB/308 ⊠ Form PCT/ISA/210 Frenc	h version 🔀 Form PCT/ISA/210 English Version			
		•	h Version  Form PCT/IPEA/409 Transla				
1			h version Form PCT/IPEA/416 English				
	□ PCT/ FR00/02452 as published    □ Form PCT/IB/306						

## JC13 Rec'd PCT/PTO 0 4 MAR 2002

U.S. APPLICATION NO. NOT APPLICABLE	RNEYS DOCKET NU <b>)110</b>	JMBER							
24. The following fees are	CALCULATIONS	PTO ONLY							
BASIC NATIONAL FEI									
Neither international preliminary examination fee (37 CFR 1,482) nor international search fee (37 CFR 1.445(a)(2) paid to USPTO and International Search Report not prepared by the EPO or JPO									
International preliminary examination fee not paid to USPTO but International Search Report prepared by the EPO or JPO									
International prelimi	inary examination (37 CFR 1,4445 (a	fee (37 Cl a)(2)) fee p	FR 1.482) not paid to USPT paid to USPTO	O but	6740.00				
☐ International prelimi	The state of the s								
International prelimi	inary examination	fee (37 Cl	FR 1.482) paid to USPTO a -(4)	nd all	\$100.00				
·	EN	TER A	PPROPRIATE BASI	C FEE AMOU	J <b>NT</b> =	\$890.00			
Surcharge of \$130.00 for months from the earliest of				20 🔲 30		\$0.00			
CLAIMS	NUMBER FI		NUMBER EXTRA	RATE			· · · · · · · · · · · · · · · · · · ·		
Total Claims	20	- 20 =	0	X \$18.00		\$0.00			
Independent Claims	1	-3 =	0	X \$84.00		\$0.00			
Multiple Dependent Clair	ns (Check if appli					\$0.00			
			OTAL OF ABOVE			\$890.00			
Applicant claims sm	nall entity status. (S	See 37 CF	R 1.127). The fees indicate			\$0.00			
				SUBTO		\$890.00			
Processing fee of \$130.00 earliest claimed priority d			ranslation later than 20	30 months fi	om the	\$0.00			
			TOTAL	NATIONAL	FEE =	\$890.00			
Fee for recording the enclaceompanied by an appro	losed assignment ( priate cover sheet	37 CFR 1 (37 CFR )	.21(h)). The assignment mu 3.28, 3.31) (check if applic	st be [able)		\$0.00	L		
			TOTAL	FEES ENSLO	SED =	\$890.00			
						Amount to be: refunded	\$		
_						charged	\$		
<del>-</del>			cover the above fees is end 3-1171 in the amount of \$		he above	fees.			
c.  The Commiss	ioner is hereby aut	thorized to	charge any additional fees				nt to Deposit		
Account No. 1	charged to a credi	t card. W	f this sheet is enclosed. ARNING: Information on t	his form may beco	me public	c. Credit card inform	ation should not		
NOTE: Where an appro	opriate time limit	under 37	card information and author CFR 1.494 or 1.495 has n			evive (37 CFR 1.137	(a) or (b)) must be		
filed and granted to res	tore the application	on to pend	ding status.		γ ·	C.M.			
SEND ALL CORRESPO	NDANCE TO			Smr (	<u> </u>	- uccu	4		
John A. SHEDDEN	1		SIC	NATURE					
RHODIA INC.			Jo	hn A. SHEDD	EN				
259 Prospect Plains	s Road		NA	ME					
CN 7500			25	,644					
CRANBURY, NJ	8512		RE	GISTATION NUM	1BER				
TEL: (609) 860-4190			M	ARCH 1	, 2002				
			DA	TE					



#### Case RN99110

#### IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In the Application of: Olivier NORE, and Jean-Luc SIMON

Examiner: N/A National Phase of PCT/FR00/02452

International Filing Date: September 6, 2000

Serial No: To be assigned

Art Unit: N/A

Filing Date: To be assigned

For: PROCESS FOR PRODUCTION OF EXOPOLYSACCHARIDES

Commissioner for Patents Washington, D.C. 20231

#### PRELIMINARY AMENDMENT

Dear Sir:

Prior to calculation of filing fee, please enter the following amendment in the specification and claims:

#### In the Specification:

Page 1, just after the title, please add the new following paragraph:

This application is an application under 35 U.S.C. Section 371 of International Application Number PCT/FR00/02452 filed on September 6, 2000.

#### In the Claims:

Please cancel claims 1-24 and replace them with the following new claims 17-36:

17. (New) A process for producing exopolysaccharides comprising a fermentation of microorganisms step, wherein the fermentation is carried out in a nutrient medium containing at least one carbon source assimilable by the microorganisms and at least one organic nitrogen source, said source deriving from a fraction of a carob bean.

Ť

- 18. (New) The process according to claim 1, wherein the fraction of the carob bean has a protein content of at least 45% by weight with respect to the dry weight of the dry matter.
- 19. (New) The process according to claim 18, wherein the protein content is of at least 60%.
- 20. (New) The process according to claim 18, wherein the protein has a high content of arginine, of glutamine or glutamic acid, and of lysine.
- 21. (New) The process according to claim 17, wherein the fraction of the carob bean has a content of lipids of at least 4% by weight with respect to the dry matter.
- 22. (New) The process according to claim 21, wherein the content of lipids is of between 7 and 15%
- 23. (New) The process according to claim 17, wherein the fraction is a germ of the carob bean.
- 24. (New) The process according to claim 17, wherein the carob bean fraction is in the form of a flour.
- 25. (New) The process according to claim 24, wherein the flour has a granulometry of between 10 and 150 microns.
- 26. (New) The process according to claim 17, wherein the nutrient medium further contains at least one inorganic nitrogen source.
- 27. (New) The process according to claim 26, wherein the inorganic nitrogen source is an ammonium or sodium nitrate, an ammonium phosphate or sulfate, a magnesium sulfate, a potassium or sodium sulfate, or a mixture thereof.

- 28. (New) The process according to claim 17, wherein the organic and optionally inorganic nitrogen source in the fermentation medium is in a concentration of between 1 and 80 g/l.
- 29. (New) The process according to claim 28, wherein the concentration of organic and optionally inorganic nitrogen source is of between 5 and 30 g/l.
- 30. (New) The process according to claim 17, wherein the assimilable carbon source is a glucose or a sucrose.
- 31. (New) The process according to claim 17, wherein the assimilable carbon source is in a concentration of between 1 and 100 g/l.
- 32. (New) The process according to claim 31, wherein the concentration of assimilable carbon source is of between 15 and 80 g/.
- 33. (New) The process according to claim 17, wherein fermentation of the microorganisms is carried out without an enzyme.
- 34. (New) The process according to claim 17, wherein fermentation is carried out at a temperature of between 15 and 100°C.
- 35. (New) The process according to claim 34, wherein the temperature is of between 25 and 35°C.
- 36. (New) The process according to claim 17, wherein the microorganism is selected from the group consisting of bacterias of the genus *Xanthomonas*, bacterias of the genus *Alcaligenes*, bacterias of the genus *Agrobacterium*, bacterias of the genus *Arthrobacter*, bacterias of the genus *Azotobacter*, bacterias of the genus *Pseudomonas*,

bacterias of the genus *Corynebacterium*, bacterias of fungi of the genus *Sclerotium*, bacterias of the genus *Aspergillus*, and yeasts of the genus *Hansenula*.

#### **REMARKS**

The preliminary amendments are filed to comply with the claims structure and wording according to the United States Patent law. It is asserted that these amendments do not add new matter. Support for these amendments can be found in the specification and claims as originally filed.

New claims find basis as mentioned in the chart below:

New Claims	Basis			
!	In the claims as filed	In the specification		
17	1			
18	2			
19	2			
20	3			
21	4			
22	4			
23	5			
24	6			
25	7			
26	8			
27	9			
28	10			
29	10			
30	11			
31	12			
32	12			
33	13			
34	14			
35	14			
36	15			

Entry of these amendments is respectfully requested.

Respectfully submitted,

Rhodia Inc.

259 Prospect Plains Road

march, 2002

CN7500

Cranbury, NJ 08512

RN99110 prelim doc

John A. SHEDDEN

Registration No. 25,644

Tel: (609) 860-4190 Fax: (609) 860-0503

#### DOCKET NO. RN99110

### EL822430009US

PCT/FR00/02452

1

#### PROCESS FOR PRODUCTION OF EXOPOLYSACCHARIDES

The present invention relates to a process

for the production of exopolysaccharides by
fermentation by means of microorganisms. More

5 specifically, the invention relates to a process for
the production of exopolysaccharides by fermentation of
microorganisms in a nutrient medium containing at least
one carbon source assimilable by the microorganisms and
at least one organic nitrogen source deriving from a

1 leguminous plant having a high content of proteins.

In the context of the present invention, the term exopolysaccharide denotes the polysaccharides produced by microorganisms.

The exopolysaccharides of high molecular

weight are increasingly used in numerous industrial

applications for their thickening, viscosifying,

emulsifying, stabilizing properties in especially

aqueous media. Thus, xanthan gum, because of its

exceptional rheological properties, is used in areas as

varied as the building industry, painting, paper, the

textile industry, cosmetics, food, agriculture, water

treatment, drilling, oil recovery etc.

These exopolysaccharides have high molecular weights, most often greater than 1 x 10<sup>6</sup> g/mol (measured by gel permeation), and are formed of units of glucose, mannose, galactose, rhamnose, glucuronic acid,

mannuronic acid, guluronic acid, optionally with acetate and pyruvate derivatives. Their particular structure and their properties are described, for example, in the work Industrial Gums - Whistler - 2nd 5 Edition - Chapters XXI-XXIII (1973).

The exopolysaccharides are advantageously produced by aerobic culture of microorganisms in an aqueous nutrient medium.

Xanthan gum is produced by bacteria of the
10 genus Xanthomonas. The exopolysaccharides of the same
type can be produced by a great variety of
microorganisms including, among the most well known,
those of the genus Agrobacterium, Arthrobacter,
Alcaligenes (Succinoglycan), Pseudomonas (Levan),
15 Rhizobium, Sclerotium (Scleroglucan).

The aqueous nutrient medium normally comprises, apart from various growth elements, a carbon source and a nitrogen source. In the industrial fermentations, the choice of the carbon source and/or of the nitrogen source is at the same time based on its availability, on its cost and on its ability to allow high productivities.

20

In certain industries, such as, for example, the food or cosmetic industry, additional constraints operate. In these areas, the carbon and nitrogen sources must, in addition, be chosen so as to obtain

3

exopolysaccharides satisfying the organoleptic, sensory and visual requirements sought.

Among the carbon and/or nitrogen sources customarily used, it is not easy to find sources which at the same time meet all the abovementioned requirements.

For example, in the case where the microorganism is not capable of consuming all of the nitrogen source, insoluble residual products remain at the end of fermentation which on the one hand make the 10 medium favorable to the development of contaminant strains which are able to degrade the must before separation of the exopolysaccharide, and on the other hand risk coloring the exopolysaccharide during 15 possible sterilization and clarification heat treatments. In certain fermentation processes, to remedy this disadvantage, it is proposed to use enzymes. Others employ filtration and/or centrifugation steps. Whatever the process of elimination of insoluble 20 residual products at the end of fermentation, an increased cost of production results therefrom.

Certain carbon and/or nitrogen sources have the disadvantage of considerably prolonging the fermentation cycle involving, especially, the contamination and thus the degradation of the must

25

4

before separation of the exopolysaccharide, and the loss of production.

The nature of the nitrogen source is particularly important when it is sought to obtain an exopolysaccharide having good organoleptic, sensory and visual properties. It is also responsible for the good productivity of the exopolysaccharide.

It has been noted that certain sources deriving from a fraction of the seed of certain

10 leguminous plants, such as the carob bean, was an organic nitrogen source of particular interest in the fermentation of microorganisms. These fractions have proved to satisfy all of the abovementioned requirements.

- Dean, those advantageously having a high content of proteins give particularly interesting results, especially in terms of productivity. On "standard" media such as those mentioned, for example, in the work Industrial Gums Whistler 2nd Edition Chapters XXI-XXIII (1973), for control fermentations, the productivities are of the order of 0.3 to 0.4 g/(kg.h); for fractions deriving from the carob bean, this productivity is greater than 0.4 g/(kg.h).
- 25 The aim of the present invention is to propose a process for production of exopolysaccharides

5

by fermentation of microorganisms, which is simple and economical.

Another aim of the invention is to propose a process for production of exopolysaccharides by

5 fermentation of microorganisms which avoids the problems of contamination explained above.

Thus, the invention relates to a process for production of exopolysaccharides by fermentation of microorganisms, characterized in that the fermentation is carried out in a nutrient medium containing at least one carbon source assimilable by the microorganisms and at least one organic nitrogen source, said source deriving from a fraction of the carob bean.

10

Other advantages linked to the choice,

15 especially, of the nitrogen source, are the reduction
of the time of the fermentations, the suppression of
insoluble residual products at the end of fermentation
and an improved productivity.

In addition, this process allows an 20 exopolysaccharide to be obtained having good organoleptic, sensory and visual properties.

Furthermore, the rheological properties of the exopolysaccharide obtained by this process are preserved and even improved in certain cases.

25 The process of the invention is capable of being applied to the production of any exopolysaccharide by

fermentation by means of microorganisms. Numerous microorganisms such as bacteria, yeasts, fungi, algae, are capable of producing exopolysaccharides. It is possible to mention, among others:

- bacteria belonging to the genus Xanthomonas and more particularly to species described in Bergey's Manual of Determinative Bacteriology (8th edition 1974 -Williams N. Wilkins Co. Baltimore) such as Xanthomonas begoniae, Xanthomonas campestris,
- 10 Xanthomonas carotae, Xanthomonas hederae, Xanthomonas incanae, Xanthomonas malvacearum, Xanthomonas papavericola, Xanthomonas phaseoli, Xanthomonas pisi, Xanthomonas vasculorum, Xanthomonas vesicatoria, Xanthomonas vitians, Xanthomonas pelargonii;
- bacteria belonging to the genus Arthrobacter and more particularly the species Arthrobacter stabilis, Arthrobacter viscosus;
  - bacteria belonging to the genus Erwinina;

20

25

- bacteria belonging to the genus Azotobacter and more particularly the species Azotobacter indicus;
- bacteria to the genus Agrobacterium and more particularly the species Agrobacterium radiobacter, Agrobacterium rhizogenes, Agrobacterium tumefaciens;
- bacteria belonging to the genus Alcaligenes and more particularly Alcaligenes faecalis;

- bacteria belonging to the genus Pseudomonas and more particularly Pseudomonas methanica;
- bacteria belonging to the genus Corynebacterium;
- bacteria belonging to the genus Bacillus and more particularly Bacillus polymyxa;

5

- fungi belonging to the genus Sclerotium and more particularly to the species Sclerotium glucanicum, Sclerotium rolfsii or Plectania occidentalis;
- fungi belonging to the genus Aspergillus and more
   particularly to the species Aspergillus itaconicus,
   Aspergillus terreus;
  - yeasts belonging to the genus *Hansenula* such as the species *Hansenula capsulata*.

Preferably, the microorganism is a bacterium of the genus Xanthomonas and more particularly of the species Xanthomonas campestris.

The invention principally relates to a process for production of exopolysaccharides by fermentation of microorganisms, characterized in that the fermentation is carried out in a nutrient medium containing at least one carbon source assimilable by the microorganisms and at least one organic nitrogen source, said source being derived from a fraction of the carob bean.

25 The carob tree produces a fruit formed of two parts, the pod and the bean. The carob bean, and more

particularly the endosperm fraction of this bean, is already widely developed under the name "carob bean gum". Closely related to this endosperm fraction is the germ, which is a by-product obtained in large 5 quantities during the isolation of the carob bean gum.

Among the different fractions of the carob bean, it turns out that all those having a high content of proteins were more particularly suited to the process of the present invention.

- Thus, the fraction of the carob bean advantageously has a protein content of at least 45%, preferably of at least 50%, and more preferentially of at least 60%, by weight with respect to the dry weight of the dry material.
- The protein content is calculated from the measurement of the nitrogen liberated by combustion at 950°C under oxygen and measured by conductivity in a stream of helium. The apparatus used is a LECO FP 428.

These proteins are formed as much of 20 essential amino acids as of nonessential amino acids.

25

A particular embodiment of the invention consists in employing fractions of the carob beans whose proteins advantageously have a high content of arginine, of glutamine and/or of glutamic acid, and of lysine.

9

In this particular embodiment, the arginine content is advantageously between 9 and 20%, and preferably between 12 and 14%, weight/weight with respect to the total of amino acids.

In the same fashion, the content of glutamine and/or glutamic acid is advantageously between 18 and 30%, preferably between 22 and 27%, weight/weight with respect to the total of amino acids.

The content of lysine is advantageously

10 between 18 and 30%, preferably between 12 and 14%,

weight/weight with respect to the total of amino acids.

The content of amino acids is determined by methods which are conventional and known to the person skilled in the art.

15 Apart from the proteins, the fractions can likewise contain lipids. The exopolysaccharides produced by fermentation of microorganisms in a nutrient medium containing at least one organic nitrogen source derived from a fraction of the carob 20 bean containing lipids more particularly see their organoleptic, visual and sensory properties markedly improved. These lipids likewise prevent the foaming in the preculture phases.

Advantageously, the content of lipids in said 25 fractions is at least 4%, advantageously at least 5%,

10

and even more advantageously varies between 7 and 15% by weight with respect to the dry matter.

The content of lipids is reduced to that of the total dry matter. It is determined by extraction with hexane in a Soxhlet extractor. The operating mode is as follows:

- exactly about 10 g of carob germ flour, say E grams, are weighed into the cartridge of the extractor, sealed with a plug of absorbent cotton;
- 10 150 ml of hexane are introduced into a 250 ml roundbottomed flask, previously tared (P0 grams);
  - the mixture is extracted for about 6 hours;

15

- the solvent is evaporated using a rotary evaporator and the drying of the residue is completed in an oven at 105°C for 1 hour;
- after cooling in a desiccator, the round-bottomed flask containing the residue, say P1 grams, is weighed.

The content of fatty matter and thus of lipids is

determined according to the following formula:

Content of fatty matter (%) =  $100 \times (P1 - P0)/E$ 

Among the characteristic compounds present in these lipids, it is especially possible to mention palmitic, stearic, oleic and linoleic acids.

11

Beside the proteins and the lipids, the fractions of the carob bean can also contain carbohydrates.

According to a particular embodiment of the invention, said fraction is the germ of the carob bean.

In this embodiment, said fraction is first freed of its endosperm fractions according to the known conventional methods.

The fraction of the carob bean used can preferably be in the form of a flour. The flour is obtained by conventional means of grinding such as grinding in mills of the type:

- cylinder mills for flours of average granulometry 100 mesh type, that is to say a flour having at most 1% by
- 15 mass of particles of greater than 80 mesh and at most 10% by mass of particles of less than 200 mesh:
  - pin mills for flours of finer granulometry:

20

25

- 200 mesh type, that is to say a flour not having particles of greater than 80 mesh and having at most 60% by mass of particles of less than 200 mesh, and
- 175 mesh type, that is to say a flour having at most 1% by mass of particles of greater than 80 mesh and at most 75% by mass of particles of less than 200 mesh.

The flour can be used as it is or after treatment by suitable enzymes such as, for example, alkaline, acid and/or neutral proteases; lipases; phytases; alkaline, acid and/or neutral phosphatases; amylases. The treatment by the enzymes is carried out by conventional and known methods.

The granulometry of said flour can fluctuate between 10 and 150 microns. In the case of treated flours, this granulometry is more particularly from 20 to 60 microns, preferably from 30 to 50 microns.

The granulometry measurements can be carried out by the laser granulometry technique, with the aid of a MALVERN granulometer, marketed by Malvern Instruments S.A.

It is likewise possible to envisage the use of the fraction of the carob bean as it is, that is to say after separation of the endosperm in the form of platelets, or even in the form of an aqueous predispersion or presuspension.

15

Although the invention is described for the carob bean, it can likewise apply to other leguminous plants such as, for example, guar, cassia, tara. These leguminous plants are mentioned in an indicative and nonlimiting capacity.

In another particular embodiment of the invention, the fermentation takes place with a mixture of organic and inorganic nitrogen sources.

In this case, the inorganic nitrogen source can be chosen from the ammonium or sodium nitrates, the ammonium phosphates or sulfates, magnesium sulfate, potassium or sodium sulfate, on their own or as a mixture.

The concentration of organic and optionally inorganic nitrogen sources in the fermentation medium is between 1 and 80 g/l, preferably between 3 and 50 g/l, and more preferentially between 5 and 30 g/l.

The fermentation medium likewise contains a source of carbon assimilable by the microorganisms.

- As constitutive carbon source of the fermentation medium, it is possible to mention glucose, sucrose, fructose, galactose, trehalose, mannose, melobiose, raffinose, maltotriose, maltose, lactose, lactulose, methyl- $\beta$ -galactopyranoside, methyl- $\alpha$ -
- galactopyranoside, cellobiose, gentobiose, methyl-β-Dglucopyranoside, methyl-α-D-glucopyranoside, esculin,
  ribose, arabinose, xylose, palatinose, rhamnose,
  fucose, melezitose, D(+) arabitol, L(-) arabitol,
  xylitol, dulcitol, tagatose, glycerol, myoinositol,
- 25 mannitol, maltitol, turanose, sorbitol, adonitol, lyxose, erythritol advantageously hydrolyzed starch,

starch hydrolyzates, mixtures of these sugars, and mixtures comprising at least one of these sugars.

Glucose and sucrose are the preferred sugars.

The concentration of assimilable carbon source is between 1 and 100 g/l, and preferably between 15 and 80 g/l.

The fermentation medium can, in addition, contain oligoelements such as traces of mineral salts, such as sulfates, chlorides of iron, of calcium, of manganese, of magnesium, of sodium, of potassium, of nickel, of cobalt, of copper. of zinc or a mixture thereof, as well as vitamins, nucleotides and/or other conventional additives such as pH control agents and antifoam agents.

- The process for production of
  exopolysaccharides according to the invention by
  fermentation of microorganisms can optionally be
  carried out in the presence of enzyme(s) such as
  alkaline, acid and/or neutral proteases;
- 20 polysaccharases; amidases; peptidases, amyloglucosidases, phosphatases; phytases.

However, one of the important advantages of the process according to the invention resides in the fact that it is possible to carry out the fermentation of the microorganisms in the absence of enzyme. It has quite surprisingly been noted that in the absence of enzyme, neither the time nor the productivity of the fermentation process were affected. In addition, the suppression of enzyme did not involve an accumulation of insoluble and undissolved residual products at the end of fermentation which can render the medium favorable to the development of contaminating strains which are able to degrade the must before separation of the exopolysaccharide.

The pure culture of the microorganisms can be carried out in the conventional manner. The person skilled in the art, as a function of the microorganism, will be in a position to choose the conditions, especially the temperatures and times of incubation, and the nature of the maintenance medium of said microorganism.

For the conservation of the microorganism, it is preferable to provide for at least one preculture step. Preculture is understood as meaning a step which consists in developing and in multiplying the bacterial strain, without production of exopolysaccharide.

The microorganism is introduced into the fermentation medium in known manner by itself with the aid of inocula or intermediate cultures.

The fermentation can be carried out at pressures of between 0 and 4 bar.

It is possible to carry out the fermentation at a temperature of between 15 and 100°C, preferably between 25 and 80°C, and more particularly between 25 and 35°C.

5 The pH of the fermentation medium can vary between 5 and 9, and preferably between 6 and 8. The pH can be adjusted, according to the case, with a base such as sodium hydroxide, potassium hydroxide or ammonia, or with an acid such as sulfuric acid, 10 phosphoric acid, hydrochloric acid or nitric acid.

The fermentation medium, placed in a tank or a fermentation vessel, can be advantageously subjected to stirring and to aeration. This stirring can be conducted, for example, by means of a reciprocating stirrer, a gyratory stirrer, one or more stirring moving body(ies) or a bubble column. The fermentation time is customarily greater than 30 hours, but generally between 40 and 100 hours.

The productivity is measured as a function of the quantity of exopolysaccharide produced, expressed in grams, with respect to kg of must, per hour of fermentation. With the process of the invention, an improvement in productivity of 3 to 15%, and preferably of 5 to 10%, has been observed.

25 After completion of the fermentation, the exopolysaccharide can be recovered from the must and

17

purified according to the known methods such as filtration, concentration, crystallization or extraction by solvents.

The invention likewise covers the

5 exopolysaccharides obtained or capable of being
obtained by the process. It more particularly covers
xanthan gum produced by the process of the invention.

The xanthan gum obtained according to the process of the invention, in 1% aqueous solution in

10 distilled water, has a high transparency, that is to say of the order of 70 to 95% or even of the order of 80 to 95%. The transmittance of the aqueous solution is measured by spectrophotometry at 600 nm.

The following examples illustrate the present invention without, however, limiting the scope thereof.

#### **EXAMPLES**

#### Example 1

20

This example describes the preculture phases 1 and 2 for Xanthomonas campestris.

#### Preculture 1:

25 Composition of the preculture medium:

18

-Yeast extract	3 g/l	Oxoid
-Malt extract	3 g/l	Oxoid
-Soybean peptone	5 g/l	Oxoid
-Sucrose	10 g/l	Eurosucre

5 -pH adjusted to 7 with  $H_2SO_4$ 

-qsp 1 liter with drinking water

All the constituents are dissolved in 1 liter of drinking water, homogenized by magnetic stirring and distributed into 500 ml Erlenmeyer flasks in 112 ml

10 fractions.

The preparation is autoclaved for 30 minutes at  $120 \, ^{\circ}\text{C}$ .

The strain is initially stored in the form of tubes frozen at -196°C by the process of freezing in liquid nitrogen vapors.

For liquid nitrogen freezing, a preculture is carried out on a specific medium having the following composition:

	<pre>malt extract</pre>	3 g	(obtained from Oxoid)
	• yeast extract	3 g	(Oxoid)
20	soybean peptone	5 g	(Oxoid)
	• glucose	10 g	(obtained from Prolabo)

□ spring water qsp 1 l.

For the preparation of the medium, all the ingredients are dispersed in spring water. The pH is adjusted to 6.5 with 10% H<sub>2</sub>SO<sub>4</sub>. The medium is sterilized for 20 minutes at 120°C, in an autoclave.

19

After incubation at 28°C for 24 hours on a gyratory stirrer at 220 rpm and amplitude = 50 mm, 10% by volume of pure sterile glycerol are added to the culture. The culture is then distributed into cryotubes of capacities varying from 1 ml to 10 ml, preferably from 2 ml to 4 ml.

These tubes are stored in liquid nitrogen vapor.

The preculture 1 is seeded with the aid of a cryotube previously thawed in ambient air. All or 50% of the cryotube is sterilely introduced into the 500 ml Erlenmeyer flasks, whose medium has been autoclaved and thus sterilized in the manner described above.

The medium seeded in this way is incubated

15 for 24 hours at 28°C on a gyratory stirrer at 220 rpm

and an amplitude of 50 mm.

After 24 hours' incubation, we obtain a preculture whose pH varies from 7 to 7.5, whose viscosity is between 50 and 500 mPa.s and whose 20 bacterial population of *Xanthomonas campestris* is greater than 10<sup>10</sup>/ml.

#### Preculture 2:

Preculture 1 is used to seed preculture 2.

25

Composition of the medium of preculture 2:

20

- Sucrose	10 g /l	Eurosucre		
- Carob germ flour	4 g/l	Meyhall AG		
- Na <sub>2</sub> HPO <sub>4</sub>	3 g/l	Europhos		

- Drinking water or softened water qsp 1 l

- pH adjusted with 10% sulfuric acid to 6.5 before sterilization.

All the constituents are suspended in 1 liter of drinking water and the pH is adjusted to 6.5. The complete medium is autoclaved for 30 min at 120°C after having distributed said medium into 500 ml Erlenmeyer flasks in 112 ml fractions.

These Erlenmeyer flasks are then seeded with 0.1 to 0.2 ml of preculture 1. These Erlenmeyer flasks are incubated for 24 to 30 hours at 28°C on a gyratory stirrer at 220 rpm and an amplitude of 50 mm.

After 24 to 30 hours' incubation, we obtain a preculture whose pH varies from 5.8 to 7.1, whose viscosity is between 100 and 1 000 mPa.s and whose bacterial population of Xanthomonas campestris is greater than 109/ ml.

#### Example 3

5

This example describes the preparation and the obtainment of the exopolysaccharide according to two fermentation processes, one with an organic

21

nitrogen source and the other with a mixed organic nitrogen and inorganic nitrogen source.

In this example, two "preculture" steps are involved. These steps take place in 500 ml Erlenmeyer 5 flasks, which corresponds to 100 ml of medium (see examples 1 and 2).

The production step, which corresponds to the step in the course of which the bacterial strain produces the polysaccharide, takes place in a 20 liter fermenter, of which 15 litres are useful.

#### Preculture step 1 and 2:

The preculture steps 1 and 2 are carried out in the same manner as in examples 1 and 2.

15

#### Production step:

#### Medium 1:

The last step is the step of production of exopolysaccharide.

20

The medium of fermenter 1 has the following composition:

	D	sucrose	42	g	Eurosucre
	ø	carob germ flour	6	g	Meyhall AG
25	a	$MgSO_4.7H_2O$	0.25	g	Bittersalz
	ø	Na <sub>2</sub> HPO <sub>4</sub>	2	g	Prolabo

22

- organic antifoam 0.5 ml
- drinking or softened water qsp 1 l

The preparation of the nitrogenous and the carbohydrate sources is carried out separately.

Sucrose  $\Rightarrow$  Qsp grams of glucose are dissolved in qsp 3 l of softened or drinking water in a Mariotte flask. The pH is lowered to 5.2 with 10%  $H_2SO_4$ . The solution is sterilized in a Mariotte flask for 45 minutes at 120°C in an autoclave.

Carob germ flour + salts ⇒ Qsp g of carob germ flour, 30 g of Na<sub>2</sub>HPO<sub>4</sub>, 3.75 g of MgSO<sub>4</sub>.7H<sub>2</sub>O, and 7.5 ml of antifoam are dissolved in qsp 7 l of softened water. The pH is adjusted to 6 with 10% H<sub>2</sub>SO<sub>4</sub>. This mixture is sterilized in situ for 45 minutes at 120°C.

1N sodium hydroxide ⇒ 40 g of NaOH pellets are dissolved in qsp 1 l of distilled water. The solution is sterilized in a Mariotte flask for 30 minutes at 120°C in an autoclave.

20 When all the ingredients are at 28°C, they are mixed in the fermenter. The fermenter is then inoculated with qsp of preculture 2.

The fermentation conditions in the fermenter are as follows:

25 Stirring  $\Rightarrow$  200 rpm from 0 to 20 hours of age, then 400 rpm until the end of the fermentation

23

Aeration  $\Rightarrow$  400 1/h from 0 to 18 hours then 825 1/h from 24 hours until the end of the fermentation

The temperature is regulated at 28°C.

The pH is regulated at 6.8 with 1N NaOH.

5 The pressure is atmospheric pressure.

#### Medium 2:

# Medium 2, which can be an alternative to medium 1, has the following composition:

10		Sucrose	42 g/l	(Eurosucre)
	0	NH <sub>4</sub> NO <sub>3</sub>	1.15 g/l	(Atochem)
	۵	MgSO <sub>4</sub> .7H <sub>2</sub> O	0.25 g/l	(Bittersalz)
	0	$(NH_4)_2HPO_4$	0.217 g/l	(Europhos)
	٥	Solubles of carob germ f	flour 36 g/l	(Meyhall AG)
15	0	organic antifoam	0.2 ml	
		Character man 1 1		

• softened water qsp 1 l

Sucrose  $\Rightarrow$  Qsp g of glucose are dissolved in qsp 3 l of softened water. The pH is adjusted to 5 with 10%  $H_2SO_4$ .

20 The solution is sterilized in a Mariotte flask for 30 minutes at 120°C in an autoclave.

Nitrogen + salts ⇒ 17.25 g of NH<sub>4</sub>NO<sub>3</sub>, 3.75 g of MgSO<sub>4</sub>.7H<sub>2</sub>O, 3.22 gr of (NH<sub>4</sub>)2HPO<sub>4</sub>, 525 gr of solubles of carob germ flour and 3 ml of antifoam are dissolved in 25 gsp 7 l of softened water. The pH of this solution is

adjusted to 6 with 10%  $\rm H_2SO_4$ . This mixture is sterilized in situ for 45 minutes at 120°C.

1N sodium hydroxide ⇒ 40 g of NaOH pellets are dissolved in qsp 1 l of distilled water. The solution is sterilized in a Mariotte flask for 30 minutes at 120°C in an autoclave.

The solubles of carob germ flour are prepared by dilution of flour to 6 to 15% in softened water.

This solution can be untreated or treated with

10 alkaline, acidic and/or neutral protease type enzymes; lipases; phytases; alkaline, acidic and/or neutral phosphatases; amylases; before being decanted, if desired, on a horizontal rotary decanter in order to eliminate the impurities which could interfere with the quality of the final product.

When all the ingredients are at 28°C, they are mixed in the fermenter (medium 1 or 2). The fermenter is then inoculated with qsp of preculture 2.

The fermentation conditions in fermenter 2

20 are as follows:

Stirring  $\Rightarrow$  200 rpm from 0 to 20 hours of age, then 400 rpm until the end of the fermentation

Aeration  $\Rightarrow$  400 1/h from 0 to 24 hours then 825 1/h from 24 hours until the end of the fermentation

25 The temperature is regulated at 28°C

The pH is regulated at 6.8 with 1N NaOH

25

The pressure is atmospheric pressure or a pressure which can range from 0.5 to 4 bars.

#### Fermentation results:

- 5 According to the culture medium studied, the fermentation times vary from 45 to 65 hours, the dry matter precipitable with isopropanol varies from 20 to 30 g/kg, and the yield by weight with respect to the carbon source employed varies from 50 to 70%. The
- 10 fermentation must obtained has a luminosity and a brightness never observed with any other source of nitrogen.

26

#### **CLAIMS**

- 1. A process for production of

  5 exopolysaccharides by fermentation of microorganisms,
  characterized in that the fermentation is carried out
  in a nutrient medium containing at least one carbon
  source assimilable by the microorganisms and at least
  one organic nitrogen source, said source deriving from

  0 a fraction of the carob bean.
- 2. The process as claimed in claim 1, characterized in that the fraction of the carob bean has a protein content of at least 45%, preferably of at least 50%, and more preferentially of at least 60%, by weight with respect to the dry weight of the dry matter.
  - 3. The process as claimed in claim 2, characterized in that the proteins advantageously have a high content of arginine, of glutamine and/or glutamic acid, and of lysine.

20

25

4. The process as claimed in any one of claims 1 to 3, characterized in that the fraction of the carob bean contains a content of lipids of at least 4%, advantageously of at least 5%, and even more advantageously varies between 7 and 15% by weight by weight with respect to the dry matter.

- 5. The process as claimed in any one of claims 1 to 4, characterized in that the fraction is the germ of the carob bean.
- 6. The process as claimed in any one of claims 1 to 5, characterized in that the carob bean fraction is in the form of a flour.
  - 7. The process as claimed in claim 6, characterized in that the granulometry of the flour fluctuates between 10 and 150 microns.
- 10 8. The process as claimed in any one of claims 1 to 7, characterized in that the fermentation is carried out in a nutrient medium containing at least one inorganic nitrogen source.
- 9. The process as claimed in claim 8,

  15 characterized in that the inorganic nitrogen source is chosen from the ammonium or sodium nitrates, the ammonium phosphates or sulfates, magnesium sulfate, potassium or sodium sulfate, on their own or as a mixture.
- 20 10. The process as claimed in any one of claims 1 to 9, characterized in that the concentration of organic and optionally inorganic nitrogen source in the fermentation medium is between 1 and 80 g/l, preferably between 3 and 50 g/l, and more

preferentially between 5 and 30 g/l.

- 11. The process as claimed in any one of claims 1 to 10, characterized in that the assimilable carbon source is chosen from glucose or sucrose.
- 12. The process as claimed in any one of claims 1 to 11, characterized in that the concentration of assimilable carbon source is between 1 and 100 g/l, and preferably between 15 and 80 g/.
- 13. The process as claimed in any one of claims 1 to 12, characterized in the fermentation of the microorganisms is carried out in the absence of enzyme.
- 14. The process as claimed in any one of claims 1 to 13, characterized in that the fermentation is carried out at a temperature of between 15 and 100°C, preferably between 25 and 80°C, and more particularly between 25 and 35°C.
- 13. The process as claimed in any one of claims 1 to 14, characterized in that the microorganism is chosen in the group of bacteria of the genus

  20 Xanthomonas, of the genus Alcaligenes, of the genus Agrobacterium, of the genus Arthrobacter, of the genus Azotobacter, of the genus Pseudomonas, of the genus Corynebacterium, of fungi of the genus Sclerotium, of the genus Aspergillus, and of yeasts of the genus

  25 Hansenula.

29

16. An exopolysaccharide obtained by a process as claimed in any one of claims 1 to 15.

#### PROCESS FOR PRODUCTION OF EXOPOLYSACCHARIDES

#### Abstract of the Disclosure

The invention concerns a method for producing exopolysaccharides by fermenting micro-organisms characterised in that it consists in carrying out the fermentation in a nutrient medium comprising at least a source of carbon available to the micro-organisms and at least a source of nitrogen, said source being derived from a fraction of carob seed.

F/Brevet/RN99110.Abst.doc

Express Mail Label #EL8	22430009US			DOGUET NO.				
		AND POWER OF ATTORNEY	ATTORNEY'S RN99110	DOCKET NO				
(include Reference to PCT Inter	rnational Applications) PC17	FR00/02452	KN99110					
As a below named inventor, I h	nereby declare that.							
My residence, post office address and citizenship are as stated below next to my name.								
I believe I am the original, first are listed below) of the subject	and sole inventor (if only one matter which is claimed and f	name is listed below) or an origi for which a patent is sought on th	nal, first and joint in ne invention entitle	nventor (if plural names d				
PROCESS FOR PRODUCTION OF EXOPOLYSACCHARIDES								
the specification of which (che	ck only one item below):							
is attached heret	o.							
was filed as Unit	ed States application							
Serial No.								
on								
and was amende	ed							
on		(if applicable)						
⊠ was filed as PC	Finternational application							
Number PCT/	FR00/02452 /		<u> </u>					
on <b>Septem</b>	ber 06, 2000,							
and amended u	nder PCT ARTICLE 19							
on	(if applicable).							
by any amendment referred to	o above.	ents of the above-identified spec						
Code of Federal Regulations.	§1.56(a).							
I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:								
PRIOR FOREIGN/PCT APP	LICATION(S) AND ANY PRIO	RITY CLAIMS UNDER 35 U.S.C	), 119:					
COUNTRY PCT indicate PCT	APPLICATION NUMBER	DATE OF FILING		ITY CLAIMED				
FRANCE _	99/11176 -	07 September, 1999	X YES	□ NO				
TIONIOL _			YES	□ NO				
			YES	□ NO				
			YES	□ NO				
			YES	□ NO				

PTO 1390 (REV 10/83)

PAGE 1 of 2

US DEPARTMENT OF COMMERCE Patent and Trademark

Office

C/Decl/RN99110 JAS Dec2

	COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY  (Continued) (include Reference to PCT International Applications) PCT/FR00/02452 / RN99110									
	I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application.									
	PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C., 120									
t			J.S. APPLICAT			US (CHECK ON				
-	υs	NUMBER	U.S FILING DATE		PATENTED		PENDING ABANDONED			
}	PCT	APPLICATIONS D	ESIGNATING	THE U.S.						
İ			T FILING DATE	US SERIAL NUMBE ASSIGNED (if any						
3-	numbe	application and transact all business in the Patent and Trademark Office connected therewith (List name and registration number) JOHN A. SHEDDEN - Reg. No. 25.644,  JOHN D. WOOD - Reg. No. 31,146  Kevin McVEIGH - Reg. No. 33,017								
. }	Send Correspondence to:					Direct Telephone Calls to:				
1	JOHN	A. SHEDDEN		_		JOHN A. SHEDDEN				
l		LECTUAL PRO	PERTY DEP	L	(609) 860-4190					
1	RHODIA INC (609) 860-4190 259 PROSPECT PLAINS ROAD,									
	CN 7500, CRANBURY, NJ 08512-7500				FIRST GIVEN		SECOND GIVEN	NAME		
-00	201	201 FULL NAME FAMILY NAME OF INVENTOR NORE			Olivier					
		RESIDENCE & CITIZENSHIP	AZAY-SU	R-CHER FX	STATE OR FOREIGN COUNTRY FRANCE		FRANCE			
		POST OFFICE ADDRESS	POST OFFICE AI	DDRESS Serraults	AZAY-SUR-CHI	SUR-CHER		STATE & ZIP CODE/COUNTRY F-37270, FRANCE		
	202	FULL NAME	FAMILY NAME		FIRST GIVEN  Jean-Luc		SECOND GIVEN	NAME		
2-04		OF INVENTOR RESIDENCE & CITIZENSHIP	SIMON CITY LILLE	FRX	STATE OR FOREIGN COUN	ITRY	COUNTRY OF CITIZENSHIP FRANCE			
		POST OFFICE	POST OFFICE A	DDRESS	LILLE		STATE & ZIP CODE/COUNTRY F-59000, FRANCE			
	203	FULL NAME	266Rue S	Solietillo	FIRST GIVEN		SECOND GIVEN			
		OF INVENTOR  RESIDENCE & CITY  CITIZENSHIP  STATE OR FOREIGN COUNTRY			ITRY	COUNTRY OF CITIZENSHIP				
		POST OFFICE	POST OFFICE A		CITY		STATE & ZIP CODE/COUNTRY			
	I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.							אונס מווט נווכ וותכ סט		
	SIGN.	Statements may jed ATURE OF INVEN	TOR 201	SIGNATURE OF	NVENTOR 202	SIGNATURE	E OF INVENTOR 203			
-	DATE			DATE JUNE	2th, 1002	DATE				
	PTO	1391 (REV 10 83	<del> </del>		SE 2 OF 2					